Commentary

QJM

Is atherosclerosis caused by high cholesterol?

U. RAVNSKOV

Introduction

According to the low-density-lipoprotein (LDL) receptor hypothesis, development of atherosclerosis is caused by a high concentration of LDL-cholesterol in the blood, and lowering LDL-cholesterol reverses, or at least retards, atherosclerosis, thus preventing cardiovascular disease.¹ As a scientific hypothesis, it is open to falsification: if the concentration of LDL-cholesterol or total cholesterol and the degree of atherosclerosis do not correlate, or if there is no exposure-response, e.g. if there is no association between the cholesterol changes (Δ LDL-cholesterol or Δ total cholesterol) and atherosclerosis progression.

The successful statin trials, with their substantial reduction of LDL-cholesterol seemed to confirm the LDL receptor hypothesis, but their outcome was independent of the initial cholesterol concentration and the degree of its lowering. For instance, the *p* values for the relationships between the outcome, and the percentage or the absolute change in LDL cholesterol, as calculated in one of the trial reports,² were 0.76 and 0.97, respectively. The lack of exposure-response, together with the benefit of the treatment in disorders and age groups where LDL-cholesterol concentration has little if any predictive value, suggests that statins must have more important effects on cardiovascular disease than a lowering of cholesterol.³ Indeed, there is evidence that the statins have anti-thrombotic and anti-inflammatory effects, and also a beneficial influence on endothelial dysfunction, LDL oxidation, re-vascularization and smooth muscle cell proliferation.

Even if these effects were operating in the trials, the substantial lowering of LDL-cholesterol

should at least have contributed to the improvement if the LDL receptor hypothesis were correct. The lack of exposure-response also questions whether atherosclerosis is truly caused by high LDL-cholesterol.

However, the outcome in the clinical trials was cardiovascular disease, not atherosclerotic progression. To answer the question, we need to compare the cholesterol concentration and the degree of atherosclerosis, and in particular, to study the influence of Δ LDL-cholesterol on atherosclerotic progression, rather than clinical outcome.

Cholesterol does not predict degree of atherosclerosis at autopsy

In 1936, Landé and Sperry noted that the degree of aortic atherosclerosis at autopsy of healthy individuals who had died violently, was independent on their blood cholesterol concentration analysed immediately after death.⁴ Their finding was confirmed by Mathur *et al.*⁵ and similar results were obtained by others.⁶⁻⁸ The objection that an analysis of cholesterol after death may not reflect its concentration during life was met by Mathur et al.⁵ who found that the cholesterol concentration was almost constant up to 16 h after death. Paterson et al.⁶ bypassed the problem by comparing the degree of atherosclerosis at death with the individuals' cholesterol measured previously on several occasions. In all these studies, plots of blood cholesterol concentrations vs. the lipid content of the aorta or the coronary arteries were widely scattered.

Address correspondence to Dr U. Ravnskov, Magle Stora Kyrkogata 9, S-22350 Lund, Sweden. e-mail: uffe.ravnskov@swipnet.se

More recent autopsy studies have found weak or inconsistent correlations between LDL-cholesterol or total cholesterol and various measures of atherosclerosis.⁹ For instance, the most severe degree of atherosclerosis was found mainly in individuals with extremely high cholesterol, whereas small differences were seen in the rest.¹⁰ A correlation was found in White men, but not in Black men,¹¹ in men but not in women,¹² in individuals below, but not above age 80 years,¹³ and in the coronary arteries, but not in the thoracic or abdominal aorta.¹⁴

The weak and unpredictable correlations probably reflect bias, because most of the studies were performed on selected individuals. In such large projects, the main object of which was to study risk factors for cardiovascular disease, individuals with such diseases, or with high cholesterol, were preferred for post-mortem examination,^{10–15} which means that the proportion of individuals with familial hypercholesterolaemia must have been much larger than in the general population. As such patients have very high cholesterol and are more prone to vascular changes, their inclusion automatically creates a correlation between degree of atherosclerosis and LDL or total cholesterol. Accordingly, it is obvious from a figure in a preliminary report that the correlation disappears if individuals with total cholesterol >350 mg/ml (9 mmol/l) are excluded.¹⁶ It is questionable if the vascular changes seen in familial hypercholesterolaemia are synonymous with atherosclerosis.17,18 Therefore, to prove that the concentration of LDL-cholesterol has importance in the general population, it is necessary to exclude individuals with familial hypercholesterolaemia.

Cholesterol does not correlate with degree of coronary atherosclerosis on angiography

A correlation between the pathological findings seen on coronary angiography and cholesterol has been found in many studies.¹⁹ However, the correlation coefficients in these studies were never > 0.36 and often much smaller; in some studies no correlation was found.^{20–23} When present, the correlation found may have been due to bias by the process mentioned above, because coronary angiography is mainly performed on patients with symptomatic coronary disease, and more often on middle-aged and younger patients. The correlation disappeared in one study after exclusion of patients treated with lipid-lowering drugs.²⁴

Cholesterol does not correlate with degree of coronary calcification

In contrast to conventional angiography, electron beam angiography detects coronary plaques independent of their location in the vessel wall, but only calcified plaques. Degree of coronary calcification seems a good surrogate for degree of coronary atherosclerosis, because it correlates strongly with total plaque volume and obstructive coronary disease, and is a powerful predictor of clinical outcome. Nonetheless, degree of coronary calcification did not correlate with any lipid fraction in the blood.²⁵

Cholesterol does not correlate with degree of peripheral atherosclerosis

Many studies have found an association between LDL- or total cholesterol and peripheral atherosclerosis, depicted by angiography or ultrasonography, but only in dichotomous analyses, and again, differences have been found mainly between individuals with very high cholesterol concentrations and the rest. In ultrasonographic studies, where degree of carotic atherosclerosis was graded as a continuous variable, no correlation was found with individual LDL-cholesterol concentrations. 26,27 In similar studies using $\operatorname{aortic}^{28}$ and femoral²⁹ angiography, no correlation was found either. Mean femoral intima-media thickness was evaluated by ultrasonography in patients with familial hypercholesterolaemia and in control individuals with normal cholesterol. Using all observations, a correlation was found (r = 0.41), but from a visual judgement of the scatterplot, within each group no clear correlation was present.³⁰

No exposure-response

The lack of an association in these studies may be explained by an influence of other important risk factors. A more reliable parameter is exposureresponse. If the amount of circulating cholesterol has any importance, sequential changes of its concentrations should be followed by parallel changes of atherosclerosis growth.

In a few observational studies with coronary angiography, the correlation of these two parameters, graded as continuous variables, was analysed. In three studies, no correlation was found;^{32–34} in two others, progression of atherosclerosis was associated with a decrease in cholesterol, not an increase.^{35,36} Experimentally, many trials have analysed the effect of cholesterol lowering on the angiographic changes. Most of them have looked at the association with on-trial LDL-cholesterol or final LDL-cholesterol only, but in sixteen trials,^{36–51} exposure-response was also analysed (Table 1). Two of them found exposure-response. In one of them Δ LDL-cholesterol and Δ total cholesterol were larger in the non-progression group, but only in a unifactorial analysis.⁴³ In another trial, treadmill exercise was used as intervention only. After one year, degree of exercise and Δ LDL-cholesterol, but not Δ total cholesterol, were inversely associated with the rate of progression.⁴⁰ In the rest of the trials exposure-response was absent (Table 1).

Several explanations were offered: most commonly that other lipids or lipid combinations explained the findings. However, Δ high-density lipoprotein (HDL) cholesterol was analysed in twelve studies, ${}^{36-38,40-44,46,47,50,51}_{47,50,51}$ Δ triglycerides in ten, ${}^{36-38,40-42,44,47,50,51}_{40,50,51}$ Δ apo-lipoprotein B in six, 37,42,47,48,50,51 Δ apo-lipoprotein A1 in three, 37,47 $\Delta very$ -low-density-lipoprotein cholesterol in three, 36,50,51 and Δ small, dense LDLcholesterol in one study,⁵⁰ but none of them were associated with atherosclerosis growth. In an early trial using visual evaluation of the angiographic findings³⁶ Δ intermediate-density lipoprotein cholesterol was associated with atherosclerotic progression, but in two others using computerassisted analysis,^{50,51} no association was found. In three trials, the ratio Δ total cholesterol/HDL cholesterol was inversely associated with athero-sclerotic progression, ^{41,43,47} but in one it was seen only in the placebo group,⁴¹ and in another the analysis was not corrected for other risk factors.⁴³

Objections

Doubt has been raised against the use of coronary angiography as a measure of atherosclerotic changes.⁵² The most serious objection, that angiography underestimates the amount of subendothelial deposits, and cannot depict the intramural ones, is not relevant in studies of exposure-response, because associations are sought to the changes, not to the degree of atherosclerosis. Large inter- and intra-observer variabilities are found in studies using visual judgement of the angiographic changes, but thirteen of the sixteen mentioned trials used quantitative, computerized image analysis (Table 1). Other objections include imprecise measurements of lumen diameter and misinterpretation of its initial compensatory enlargement as atherosclerotic regression. In particular, percent stenosis has been questioned as a reliable measure of progress or regress because of uncontrolled physiological influences on the lumen of the reference vessels.^{52–54} However, if measured with care, the minimum lumen diameter, used in half of the studies as a measure of atherosclerosis (Table 1), has a low coefficient of variation for repeated measurements⁵⁵ and is a strong predictor of the coronary flow reserve,⁵⁶ the reactive hyperaemic response,⁵³ the transstenotic pressure gradient⁵⁷ and thallium scintigraphic changes after exercise,⁵⁷ all of which reflect degree of atherosclerotic narrowing of the coronary vessels. Angiographic deterioration strongly predicted cardiovascular events in the studies that included a clinical follow-up.^{36,37,39}

Why does a high cholesterol predict cardiovascular disease?

If LDL-cholesterol and ALDL-cholesterol do not correlate with degree of atherosclerosis or with atherosclerosis growth, why does a high cholesterol predict cardiovascular disease? The answer may be that cardiovascular disease is not synonymous with atherosclerosis. A high LDL or total cholesterol may be secondary to uncontrolled factors that promote cardiovascular disease in other ways and cause hypercholesterolaemia at the same time, for instance lack of physical activity,⁵⁸ mental stress,⁵⁹ smoking, and obesity.⁶⁰ It is generally assumed that their effect on cardiovascular disease is mediated through the high cholesterol, but this may be a secondary phenomenon. Physical activity may benefit the cardiovascular system by improving endothelial function,⁶¹ or by stimulating the formation of collateral vessels;⁶² mental stress may have a harmful influence on adrenal hormone secretion, smoking increases the oxidant burden; in these all situations the high cholesterol may be an epiphenomenal indicator that something is wrong. This argument also explains why some studies found atherosclerotic growth to be associated with initial or on-study LDL-cholesterol, but not with Δ LDL or total cholesterol. If the amount of LDLcholesterol in the blood were the determining factor, atherosclerotic growth should have been associated with Δ LDL-cholesterol as well and to a higher degree.

Conclusion

'The more LDL there is in the blood, the more rapidly atherosclerosis develops.' This 1984 statement by the Nobel Award winners Michael Brown and Joseph Goldstein¹ has dominated research

	Type of intervention	Measurement of angiographic	Baseline LDL-C	Trial length (vears months)	Patien	its		Increa	e in clerosis
		progress or regress	(mmol/l)		۲	Males (%)	Mean age (years)	associa with	ited
								ΔtC	ALDL-C
Krauss <i>et al.</i> 1987 ³⁶	Cholestyramine	MLD c, v	6.08 (7.08)	5,0	143	80	I	×°N	×oN
Blankenhorn <i>et al.</i> 1990 ³⁷	Colestipol, niacin	MLD c, v	4.36 (6.26)	2,0	162	100	54	×°N	No*
Olsson <i>et al.</i> 1990 ³⁸	Nicotinic acid	Global estimate f, v	6.44 (9.68)	1,6	20	100	50	No	No
Tatami <i>et al.</i> 1992 ³⁹	+ tenotibrate LDL-apheresis	%Stenosis c. a	8.89 (11.1)	>1.0	37	59	I	*oN	*oN
	+ probucol)))
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	and/or pravastatin			0	0	00	C)	÷
Hambrecht <i>et al.</i> 1993. ⁷	Physical exercise	MLD; %stenosis c, q	4.21 (6.0)	1,0	88	100	53	° Z	Yes*
Hodis et al. 1994	Lovastatin	%stenosis c, q	4.00 (5.90)	2,0	220	91	58	* No	×0Z
Sacks et al. 1994 ⁴²	Various lipid-lowering	MLD; %stenosis c, q	3.56(5.5)	2,9	79	89	58	×°N	No*
Quinn <i>et al.</i> 1994 ⁴³	arugs Multiple risk factor	MLD c, q	4.02 (5.88)	2,0	257	86	57	Yes	Yes
:	reduction								
Schuff-Werner et al. 1994 ⁴⁴	LDL-apheresis	%Stenosis c, q	7.54 (9.36)	2,0	33	70	47	No	No
Kitabatake <i>et al.</i> 1994 ⁴⁵	LDL-apheresis + various	%Stenosis, c, q	6.62 (8.46)	1,0	13	77	48	I	No
Regnström <i>et al.</i> 1996 ⁴⁶	npia-towering arugs Probucol	Global estimate f. a	6.52 (8.83)	3.0	303	57	54	0 Z	0 N
Niebauer <i>et al.</i> 1996 ⁴⁷	Low-fat diet+physical	%Stenosis c, q	4.22 (6.06)	1,0	92	100	54	No*	No*
	exercise	-							
Kroon <i>et al.</i> 1996 ⁴⁸	LDL-apheresis + simvastatin	MLD; %stenosis c, q	7.82 (9.79)	2,0	40	100	52	No	No
Townson of $n/$ 1007 ⁴⁹	Vs. SIIIIVastaun Drovortatio	0/ Ctonocic c a	(CZ V) 11 C		00	01	73		
Runtolo <i>et al</i> 1998 ⁵⁰	Bezafihrate	MID: %stenosis r a	4 64	0, r	81	100	41		* 2 X
Sutherland <i>et al.</i> 1998 ⁵¹	Simvastatin	% stenosis c. d	4 33 (6 91)	2,0	38.		57	0 Z	o No
	טוווועמסומנווו	h 'n elentine n'		0,4			10		

400

Downloaded from by guest on September 28, 2016

on atherosclerosis since then. As shown here, this hypothesis appears to be falsified by the fact that degree of atherosclerosis, and atherosclerotic growth, were independent on the concentration or the change of LDL-cholesterol in almost all studies. The role of LDL-cholesterol for atherosclerosis growth has been exaggerated, a finding with consequences for the prevention of cardiovascular disease. For instance, as the statins exert their beneficial influence on the cardiovascular system by several mechanisms, it may be wiser to search for the lowest effective dose instead of the dose with maximal effect on LDL-cholesterol. Neither should an elevated LDL-cholesterol be the primary target in cardiovascular prevention, as recently claimed by the American National Cholesterol Education Program, and researchers should direct more attention to other hypotheses.

I may have overlooked studies that have found an association between changes of LDL-cholesterol or other lipid fractions, and atherosclerotic progression. However, although the presence of exposure-response is not sufficient proof in itself of causality, it is difficult to explain its absence.

References

- Brown M, Goldstein JL. How LDL receptors influence cholesterol and atherosclerosis. *Sci Am* 1984; 251:58–66.
- Sacks FM, Moyé LA, Davis BR, et al. Relationship between plasma LDL concentrations during treatment with pravastatin and recurrent coronary events in the cholesterol and recurrent events trial. *Circulation* 1998; **97**:1446–52.
- Ravnskov U. Implications of 4S evidence on baseline lipid levels. *Lancet* 1995; 346:181.
- Landé KE, Sperry WM. Human atherosclerosis in relation to the cholesterol content of the blood serum. *Arch Pathol* 1936; 22:301–12.
- Mathur KS, Patney NL, Kumar V, Sharma RD. Serum cholesterol and atherosclerosis in man. *Circulation* 1961; 23:847–52.
- 6. Paterson JC, Dyer L, Armstrong EC. Serum cholesterol levels in human atherosclerosis. *Can Med Ass J* 1960; **82**:6–11.
- Marek Z, Jaegermann K, Ciba T. Atherosclerosis and levels of serum cholesterol in postmortem investigations. *Am Heart J* 1962; 63:768–74.
- Cabin HS, Roberts WC. Relation of serum total cholesterol and triglyceride levels to the amount and extent of coronary artery narrowing by atherosclerotic plaque in coronary heart disease. *Am J Med* 1982; 73:227–34.
- 9. Sharrett AR. Serum cholesterol levels and atherosclerosis. *Coron Artery Dis* 1993; **4**:867–70.
- Solberg LA, Strong JP, Holme I, *et al.* Stenoses in the coronary arteries. Relation to atherosclerotic lesions, coronary heart disease, and risk factors. The Oslo Study. *Lab Invest* 1985; **53**:648–55.
- 11. Oalmann MC, Malcom GT, Toca VT, Guzman MA, Strong JP. Community pathology of atherosclerosis and coronary heart disease: post mortem serum cholesterol and extent of

coronary atherosclerosis. *Am J Epidemiol* 1981; **113**:396–403.

- 12. Feinleib M, Kannel WB, Tedeschi CG, Landau TK, Garrison RJ. The relation of antemortem characteristics to cardiovascular findings at necropsy. The Framingham Study. *Atherosclerosis* 1979; **34**:145–57.
- Sadoshima S, Kurozumi T, Tanaka K, et al. Cerebral and aortic atherosclerosis in Hisayama, Japan. Atherosclerosis 1980; 36:117–26.
- Reed DM, Strong JP, Resch J, Hayashi T. Serum lipids and lipoproteins as predictors of atherosclerosis. An autopsy study. *Arteriosclerosis* 1989; **9**:560–4.
- 15. Sorlie PD, Garcia-Palmieri MR, Costas R, Oalmann MC, Havlik R. The relation of antemortem factors to atherosclerosis at autopsy. The Puerto Rico Heart Health Program. *Am J Pathol* 1981; **103**:345–52.
- Solberg LA, Hjermann I, Helgeland A, Holme I, Leren PA, Strong JP. Association between risk factors and atherosclerotic lesions based on autopsy findings in the Oslo study: a preliminary report. In: Schettler G, Goto Y, Hata Y, Klose G, eds. *Atherosclerosis IV*. Proc 4. Int. Symp. Berlin, Springer Verlag, 1977:98–100.
- 17. Stehbens WE, Martin M. The vascular pathology of familial hypercholesterolemia. *Pathology* 1991; **23**:54–61.
- Stehbens WE. Coronary heart disease, hypercholesterolemia, and atherosclerosis I: false premises. *Exp Mol Pathol* 2001; **70**:103–19
- Pearson TA. Coronary arteriography in the study of epidemiology of coronary artery disease. *Epidemiol Rev* 1984; 6:140–66.
- 20. Nitter-Hauge S, Enge I. Relation between blood lipid levels and angiographically evaluated obstructions in coronary arteries. *Br Heart J* 1973; **35**:791–5.
- 21. Barboriak JJ, Rimm AA, Anderson AJ, Tristani FE, Walker JA, Flemma RJ. Coronary artery occlusion and blood lipids. *Am Heart J* 1974; **87**:716–21
- 22. Fuster V, Frye RL, Connolly DC, Danielson MA, Elveback LR, Kurland LT. Arteriographic patterns early in the onset of the coronary syndromes. *Br Heart J* 1975; **37**:1250–5.
- Krishnaswami S, Jose VJ, Joseph G. Lack of correlation between coronary risk factors and CAD severity. *Int J Cardiol* 1994; 47:37–43.
- 24. Cramér K, Paulin S, Werkö L. Coronary angiographic findings in correlation with age, body weight, blood pressure, serum lipids, and smoking habits. *Circulation* 1966; **33**:888–900.
- Hecht HS, Superko HR. Electron beam tomography and National Cholesterol Education Program guidelines in asymptomatic women. J Am Coll Cardiol 2001; 37:1506–11.
- Crouse JR, Toole JF, McKinney WM, et al. Risk factors for extracranial carotid artery atherosclerosis. *Stroke* 1987; 18:990–6.
- Nishino M, Sueyoshi K, Yasuno M, Yamada Y, Abe H, Hori M, Kamada T. Risk factors for carotid atherosclerosis and silent cerebral infarction in patients with coronary heart disease. *Angiology* 1993; 44:432–40.
- Palomäki H, Kaste M, Raininko R, Salonen O, Juvela S, Sarna S. Risk factors for cervical atherosclerosis in patients with transient ischemic attack or minor ischemic stroke. *Stroke* 1993; 24:970–5.

- 29. Erikson U, Ericsson M, Persson R. On the relation between peripheral atherosclerosis and serum lipoproteins. *Upsala J Med Sci* 1979; **84**:95–104.
- Wendelhag I, Wiklund O, Wikstrand J. Atherosclerotic changes in the femoral and carotid arteries in familial hypercholesterolemia. Ultrasonographic assessment of intima-media thickness and plaque occurrence. *Arteriosclerosis* 1993; 13:1404–11.
- Kramer JR, Kitazume H, Proudfit WL, Matsuda Y, Williams GW, Sones FM. Progression and regression of coronary atherosclerosis: relation to risk factors. *Am Heart J* 1983; **105**:134–44.
- 32. Bruschke AVG, Kramer JR Jr, Bal ET, Haque IU, Detrano RC, Goormastic M. The dynamics of progression of coronary atherosclerosis studied in 168 medically treated patients who underwent coronary arteriography three times. *Am Heart J* 1989; **117**:296–305.
- Bissett JK, Wyeth RP, Matts JP, Johnson JW. Plasma lipid concentrations and subsequent coronary occlusion after a first myocardial infarction. The POSCH group. *Am J Med Sci* 1993; **305**:139–44.
- 34. Bemis CE, Gorlin R, Kemp HG, Herman MV. Progression of coronary artery disease: a clinical arteriographic study. *Circulation* 1973; **47**:455–64.
- 35. Shub C, Vlietstra RE, Smith HC, Fulton RE, Elveback LR. The unpredictable progression of symptomatic coronary artery disease: a serial clinical-angiographic analysis. *Mayo Clin Proc* 1981; **56**:155–60.
- Krauss RM, Lindgren FT, Williams PT, et al. Intermediate-density lipoproteins and progression of coronary artery disease in hypercholesterolaemic men. Lancet 1987; 2:62–6.
- Blankenhorn DH, Alaupovic P, Wickham E, Chin HP, Azen SP. Prediction of angiographic change in native human coronary arteries and aortocoronary bypass grafts: lipid and nonlipid factors. *Circulation* 1990; 81:470–6.
- Olsson AG, Ruhn G, Erikson U. The effect of serum lipid regulation on the development of femoral atherosclerosis in hyperlipidaemia: a non-randomized controlled study. *J Int Med* 1990; 227:381–90.
- Tatami R, Inoue N, Itoh H, et al. Regression of coronary atherosclerosis by combined LDL-apheresis and lipid-lowering drug therapy in patients with familial hypercholesterolemia: a multicenter study. *Atherosclerosis* 1992; **95**:1–13.
- Hambrecht R, Niebauer J, Marburger C, et al. Various intensities of leisure time physical activity in patients with coronary artery disease: effects on cardiorespiratory fitness and progression of coronary atherosclerotic lesions. J Am Coll Cardiol 1993; 22:468–77.
- 41. Hodis HN, Mack WJ, Azen SP, *et al.* Triglyceride- and cholesterol-rich lipoproteins have a differential effect on mild/moderate and severe lesion progression as assessed by quantitative coronary angiography in a controlled trial of lovastatin. *Circulation* 1994; **90**:42–9.
- 42. Sacks FM, Pasternak RC, Gibson CM, Rosner B, Stone PH, for the Harvard Atherosclerosis Reversibility Project (HARP) Group. Effect on coronary atherosclerosis of decrease in plasma cholesterol concentrations in normocholesterolaemic patients. *Lancet* 1994; **344**:1182–6.
- 43. Quinn TG, Alderman EL, McMillan A, Haskell W. Development of new coronary atherosclerotic lesions during a 4-year multifactor risk reduction program: The Stanford

Coronary Risk Reduction Project (SCRIP). J Am Coll Cardiol 1994; 24:900–8.

- 44. Schuff-Werner P, Gohlke H, Bartmann U, *et al.* and the HELP-STUDY GROUP. The HELP-LDL-apheresis multicentre study, an angiographically assessed trial on the role of LDL-apheresis in the secondary prevention of coronary heart disease. II: Final evaluation of the effect of regular treatment on LDL- cholesterol plasma concentrations and the course of coronary heart disease. *Eur J Clin Invest* 1994; **24**:724–32.
- Kitabatake A, Sato H, Hori M, et al. Coronary atherosclerosis reduced in patients with familial hypercholesterolemia after intensive cholesterol lowering with low-density lipoprotein-apheresis: 1-year follow-up study. *Clin Ther* 1994; 16:416–28.
- 46. Regnström J, Walldius G, Nilsson S, *et al.* The effect of probucol on low density lipoprotein oxidation and femoral atherosclerosis. *Atherosclerosis* 1996; **125**:217–29.
- 47. Niebauer J, Hambrecht R, Velich T, *et al.* Predictive value of lipid profile for salutary coronary angiographic changes in patients on a low-fat diet and physical exercise program. *Am J Cardiol* 1996; **78**:163–7.
- Kroon AA, Aengevaeren WRM, van der Werf T, et al. LDL-apheresis atherosclerosis regression study (LAARS). Effect of aggressive versus conventional lipid lowering treatment on coronary atherosclerosis. *Circulation* 1996; 93:1826–35.
- 49. Tamura A, Mikuriya Y, Nasu M, and the coronary artery regression study (CARS) group. Effect of Pravastatin (10 mg/day) on progression of coronary atherosclerosis in patients with serum total cholesterol levels from 160 to 220 mg/dl and angiographically documented coronary disease. *Am J Cardiol* 1997; **79**:893–6.
- Ruotolo G, Ericsson CG, Tettamanti C, *et al.* Treatment effects on serum lipoprotein lipids, apolipoproteins and low density lipoprotein particle size and relationships of lipoprotein variables to progression of coronary artery disease in the bezafibrate coronary atherosclerosis intervention trial (BECAIT). *J Am Coll Cardiol* 1998; **32**:1648–56.
- Sutherland WHF, Restieaux NJ, Nye ER, et al. IDL composition and angiographically determined progression of atherosclerotic lesions during simvastatin therapy. *Arteriosclerosis* 1998; 18:577–83.
- Hong MK, Mintz GS, Popma JJ, et al. Limitations of angiography for analyzing coronary atherosclerosis progression or regression. Ann Intern Med 1994; 121:348–54.
- Harrison DG, White CW, Hiratzka LF, et al. The value of lesion cross-sectional area determined by quantitative coronary angiography in assessing the physiologic significance of proximal left anterior descending coronary arterial stenoses. *Circulation* 1984; 69:1111–19.
- White CW, Wright CB, Doty DB, et al. Does visual interpretation of the coronary arteriogram predict the physiologic importance of a coronary stenosis? N Engl J Med 1984; 310:819–24.
- 55. Ellis S, Sanders W, Goulet C, *et al.* Optimal detection of the progression of coronary artery disease: comparison of methods suitable for risk factor intervention trials. *Circulation* 1986; **74**:1235–42.
- 56. Zijlstra F, van Ommeren J, Reiber JH, Serruys PW. Does the quantitative assessment of coronary artery dimensions

predict the physiologic significance of a coronary stenosis? *Circulation* 1987; **75**:1154–61.

- 57. Wijns W, Serruys PW, Reiber JHC, *et al.* Quantitative angiography of the left anterior descending coronary artery: correlations with pressure gradient and results of exercise thallium scintigraphy. *Circulation* 1985; **71**:273–9.
- Stefanick ML, Mackey S, Sheehan M, Ellsworth N, Haskell WL, Wood PD. Effects of diet and exercise in men and postmenopausal women with low levels of HDL cholesterol and high levels of LDL-cholesterol. *N Engl J Med* 1998; 339:12–20.
- 59. Muldoon MF, Herbert TB, Patterson SM, Kameneva M, Raible R, Manuck SB. Effects of acute psychological stress

on serum lipid levels, hemoconcentration, and blood viscosity. Arch Intern Med 1995; **155**:615–20.

- 60. Ravnskov U. *The Cholesterol Myths.* Washington DC, New Trends Publishing, 2000.
- 61. Hambrecht R, Wolf A, Gielen S, *et al.* Effect of exercise on coronary endothelial function in patients with coronary artery disease. *N Engl J Med* 2000; **342**:454–60.
- Belardinelli R, Georgiou D, Ginzton L, Cianci G, Purcaro A. Effects of moderate exercise training on thallium uptake and contractile response to low-dose dobutamine of dysfunctional myocardium in patients with ischemic cardiomyopathy. *Circulation* 1998; 97:553–61.